arXiv:math/0507333v1 [math.GM] 16 Jul 2005

Recurrence Divisibility Tests

Mehdi Hassani

Department of Mathematics
Institute for Advanced Studies in Basic Sciences
Zanjan, Iran
mhassani@iasbs.ac.ir

Abstract

In this note, we are going to introduce some recurrence divisibility tests for all primes except than 2 and 5.

2000 *Mathematics Subject Classification*: 13A05, 11A41, 11B37, 11D04.
Keywords: divisibility, primes, recurrence, linear diophantine equation.

For $n \in \mathbb{N}$, we define $ud(n) =$ unite digit of n. As we know, we have the following divisibility tests for 2 and 5:

$$2|n \Leftrightarrow 2|ud(n),$$

and

$$5|n \Leftrightarrow ud(n) \in \{0, 5\}.$$

In this note, we are going to find some *recurrence divisibility tests* for other primes except than 2 and 5. As soon, you will understand our mind by "recurrence".

Theorem 1 *For every prime $p \neq 2, 5$, there exists unique $t(p) \in \mathbb{Z}_p$, in which*

$$p|n \Leftrightarrow p\Big|\left\lfloor \frac{n}{10} \right\rfloor - t(p)ud(n).$$

Also, we have

$$t(p) = \begin{cases} \frac{p-1}{10} & \text{if } ud(p) = 1, \\ \frac{7p-1}{10} & \text{if } ud(p) = 3, \\ \frac{3p-1}{10} & \text{if } ud(p) = 7, \\ \frac{9p-1}{10} & \text{if } ud(p) = 9. \end{cases}$$

Proof: Suppose $p \neq 2, 5$ is a prime. Since $\gcd(10, p) = 1$, the diophantine equation $10t + 1 = kp$ with $0 \leq t < p$, has an unique solution module p and clearly $kt > 0$. Let $t = t(p)$. So, $p|10t(p) + 1$ and $1 \leq t(p) \leq p - 1$.
Now, suppose $p|n$, or consequently $p|t(p)n$. Also, we have $p|10t(p) + 1$. Therefore, by considering $ud(n) = n - 10\lfloor \frac{n}{10} \rfloor$, we have $p|\lfloor \frac{n}{10} \rfloor - t(p)ud(n)$.
Also, suppose $p|\lfloor \frac{n}{10} \rfloor - t(p)ud(n)$ or $p|(10t(p) + 1)\lfloor \frac{n}{10} \rfloor - t(p)n$. Since $p|10t(p) + 1$, we obtain $p|t(p)n$ and since $1 \leq t(p) \leq p - 1$, we have $\gcd(p, t(p)) = 1$, therefore $p|n$.
Now, we compute the value of $t(p)$. To do this, we note that since $p \neq 2, 5$, we have $ud(p) \in \{1, 3, 7, 9\}$. If we let $k = k(p)$, we have the relation $10t(p) + 1 = k(p)p$ with $1 \leq t(p) \leq p-1$. So, $ud(pk(p)) = 1$ always holds. Also, the condition $1 \leq t(p) \leq p-1$ yield that $\frac{9}{p} \leq k(p) \leq 10 - \frac{11}{p}$. So, for all primes $p \neq 2, 5$ we have $1 \leq k(p) \leq 9$. If $ud(p) = 1$, then since $ud(pk(p)) = 1$ and $1 \leq k(p) \leq 9$, we obtain $k(p) = 1$ and so $t(p) = \frac{p-1}{10}$. Other cases have similar reasons. This completes the proof. $\square$

Note 1. We called this divisibility test, "recurrence", because for all primes $p \neq 2, 5$ we have

$$\lfloor \log_{10} n \rfloor = 1 + \left\lfloor \log_{10} \left(\left\lfloor \frac{n}{10} \right\rfloor - t(p)ud(n) \right) \right\rfloor.$$

It means, this test works by cancelling digits of given number for check, 1 digit by 1 digit.

Note 2. Same divisibility tests can be obtain which work by cancelling m digit by m digit.

Note 3. We can yield same tests for all $n \in \mathbb{N}$ with $2 \nmid n$ and $5 \nmid n$.

Acknowledgements. I deem my duty to thank Mr. H. Osanloo, who mentioned me recurrence divisibility tests in some special cases.

References

[1] G.H. Hardy and E.M. Wright, An Introduction to THE THEORY OF NUMBERS, fifth edition, Oxford University Press, London, 1979.